Trileptal (oxcarbazepine)



What is it usually prescribed for?

Partial seizures (convulsions) and bipolar disorder.



How does it work?

Trileptal helps improve seizures, thinking, mood, and behavior. Trileptal works by reducing excessive brain activity and release of brain chemicals.



What should I expect?

Trileptal is usually taken as an add-on medicine to treat bipolar disorder. It may take several weeks for you to feel the full benefits of taking this medicine. Some symptoms may improve as early as 1–2 weeks.

Your healthcare professional will start with a low dose that is slowly increased to find the best dose that works for you. Tell your healthcare professional how it's working for you and if you have any side effects, so the dose can be changed if needed.

Your healthcare professional may check your blood sodium levels, especially during the first 3 months of taking Trileptal, because this medicine can increase them.



Take Trileptal exactly as prescribed to get the full benefit of this medicine.



Will I have side effects?

You may or may not have side effects when taking Trileptal. The chance of side effects increases with larger doses.

In most cases, treatment is necessary for a long time to control symptoms. Long treatment may increase the chances of some side effects like changes in blood sodium, skin rash, and suicidal thoughts.

What are common side effects and what can I do to feel better?

It your side effects are mild, talk to your healthcare professional about them at your next appointment

	Drink plenty of water	Drink less alcohol	Do more exercise	Healthcare professional suggestion
Feeling dizzy	х	Х		
Feeling drowsy	х	Х	х	
Feeling tired		Х	х	
Nausea or throwing up	х	Х		
Feeling unsteady		Х		
Headache	х	Х	х	

Trileptal (oxcarbazepine)

YOU'RE NOT ALONE!

In 2020, Trileptal was prescribed over 4.2 million times in the United States.



What should I avoid?

Don't start any new medicines, over-the-counter drugs, or herbal remedies without talking to your healthcare professional first. There are interactions with medicines that may increase your chance of serious side effects.

Do not suddenly stop taking Trileptal unless your healthcare professional has told you to, because your symptoms may return.

Use less alcohol and recreational drugs. They may decrease the benefits of Trileptal and increase the chance of side effects.

Avoid pregnancy while taking this medicine. Trileptal is associated with birth defects and may decrease the levels and effects of oral contraceptives (birth control pills).

WHAT IS MOST IMPORTANT TO REMEMBER



It may take up to several weeks for you to feel the full benefits of taking this medicine



Tell your healthcare professional about any troubling side effects so that you can work together to address them



Do not suddenly stop taking this medicine unless told to do so by your healthcare professional



Don't start any new medicines, overthe-counter drugs, or herbal remedies without talking to your healthcare professional



Get medical care right away if you have serious side effects



When should I call a healthcare professional right away or visit the emergency room?

If you have any **serious** side effects. Here are some examples of serious side effects:

- Thoughts of suicide or are planning to hurt yourself
- Nausea, tiredness, lack of energy, headache, confusion, or more frequent or more severe seizures (convulsions)
- Skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes

If you think there has been an overdose, call your poison control center (800-222-1222) or get medical care right away.



Where can I get more help?

National Alliance on Mental Illness (NAMI) - nami.org

National Suicide Prevention Lifeline – Call 1-800-273-TALK (8255)

Crisis Text Line - Text "HELLO" to 741741

?	Notes and questions for my next appointment